

### Metadata

#### File Identifier

79afeb36-e601-20ab-b64d-807ab1b97bb0

#### Language

eng

#### Hierarchy Level Name

dataset

### Contact

#### Responsible Party

##### Individual Name

Geospatial Team

##### Organisation Name

Statistics New Zealand

##### Position Name

Geospatial Analyst

#### Contact Info

##### Contact

##### Phone

##### Telephone

##### Voice

04 931 4600

##### Address

##### Address

##### Delivery Point

Statistics House, The Boulevard, Harbour Quays

##### City

Wellington

##### Postal Code

6140

##### Country

NZ

##### Electronic Mail Address

**Date Stamp****Date Time**

20161020

**Metadata Standard Name**

ISO 19115:2003/19139

**Metadata Standard Version**

1.0

**Spatial Representation Info****Vector Spatial Representation****Integer**

0

**Identification Info****Data Identification****Citation****Citation****Title**

TA2013\_V1\_00

**Abstract**

This dataset is the definitive set of territorial authority boundaries for 2013 as defined by the Local Government Commission and/or the territorial authorities themselves but maintained by Statistics New Zealand (who is the custodian). A Territorial Authority is defined under the Local Government Act 2002 as a city or a district council. There is now a total of 67 territorial authorities in New Zealand. This total reflects the amalgamation of the seven territorial authorities (Rodney District, North Shore City, Waitakere City, Auckland City, Manukau City, Papakura District and Franklin District) into one new Auckland Council in 2010. Territorial authorities are the second tier of local government in New Zealand, below regional councils. The 67 territorial authorities comprise: 13 city councils including the Auckland council, 53 district councils, and the Chatham Islands Territory. Some territorial authority boundaries are coterminous with regional council boundaries but there are several exceptions. An example is Taupo District, which is split between four regions, although most of its area falls within the Waikato Region. When defining the boundaries of territorial authorities, the Local Government Commission based considerable weight on the 'community of interest'. Territorial authorities are defined at meshblock and area unit level.

**Purpose**

TA2013\_V1\_00 is the definitive set of territorial authority boundaries for 2013. This version contains 67 territorial authorities.

**Point Of Contact****Responsible Party****Individual Name**

Geospatial Team

**Organisation Name**

Statistics New Zealand

**Position Name**

Geospatial Analyst

**Contact Info**

**Contact**

**Phone**

**Telephone**

**Voice**

04 931 4600

**Address**

**Address**

**Delivery Point**

Statistics House, The Boulevard, Harbour Quays

**City**

Wellington

**Postal Code**

6140

**Country**

NZ

**Electronic Mail Address**

geography@stats.govt.nz

**Resource Constraints**

**Constraints**

**Use Limitation**

<DIV STYLE="text-align:Left;"><DIV><DIV><P><SPAN>These conditions of supply apply to all users of Statistics NZ digital boundaries effective 1 July 2007. Permitted uses You must acknowledge Statistics NZ as the source of the boundaries. Uses not permitted You must not change the accuracy of the boundaries and supply them to another party. Liability While care has been taken to compile these boundary coordinates, Statistics NZ gives no warranty that the data supplied is free from error. Statistics NZ will not be liable for any loss suffered by the use, directly or indirectly, of this information.</SPAN></P></DIV></DIV></DIV>

**Language**

eng

**Topic Category Code**

boundaries

## Extent

### EX \_ Extent

#### Description

Data represents subdivision polygons dissolved from meshblocks since 1991

## Extent

### EX \_ Extent

#### Geographic Element

#### EX \_ Geographic Bounding Box

165.973643757-175.379047054-47.6201235791-33.9584971002

## Distribution Info

### Distribution

#### Transfer Options

#### Digital Transfer Options

##### On Line

#### Online Resource

#### Linkage

#### URL

<https://datafinder.stats.govt.nz/layer/25735-territorial-authority-2013/>

## Data Quality Info

### DQ \_ Data Quality

#### Lineage

#### LI \_ Lineage

#### Statement

Territorial authority boundaries are based on the meshblock pattern and comprise of whole area units. Non-alignment of meshblock and cadastral boundaries are one of a number of reasons for meshblock boundary adjustments. Other reasons include requests from local authorities, Local Government Commission, Electoral Representation Commission and to make Census of Population and Dwellings enumeration processes easier. From the meshblock pattern, higher geographies, including the 2013 territorial authority pattern were dissolved using the dissolve tool in the Arc GIS suite to create multiple output datasets.

#### Source

#### LI \_ Source

#### Description

Deriving of output Files The original vertices delineating the meshblock boundary pattern were digitised in 1991 from 1:5,000 scale urban maps and 1:50,000 scale rural maps. The magnitude of error of the original digital points would have been in the range of +/- 10 metres in urban areas and +/- 25 metres in rural areas. Where meshblock boundaries coincide with cadastral boundaries the magnitude of error will be within the range of 1–5 metres in urban areas and 5 - 20 metres in rural areas. This being the estimated magnitude of error of Landonline. The creation of high definition and generalised meshblock boundaries for the 2013 digital pattern and the dissolving of these meshblocks into other

geographies/boundaries were completed within Statistics New Zealand using ESRI's ArcGIS desktop suite and the Data Interoperability extension with the following process: 1. Import data and all attribute fields into an ESRI File Geodatabase from LINZ as a shapefile 2. Run geometry checks and repairs. 3. Run Topology Checks on all data (Must Not Have Gaps, Must Not Overlap), detailed below. 4. Generalise the meshblock layers to a 1m tolerance to create generalised dataset. 5. Clip the high definition and generalised meshblock layers to the coastline using land water codes. 6. Dissolve all four meshblock datasets (clipped and unclipped, for both generalised and high definition versions) to higher geographies to create the following output data layers: Area Unit, Territorial Authorities, Regional Council, Urban Areas, Community Boards, Territorial Authority Subdivisions, Wards Constituencies and Maori Constituencies for the four datasets. 7. Complete a frequency analysis to determine that each code only has a single record. 8. Re-run topology checks for overlaps and gaps. 9. Export all created datasets into MapInfo and Shapefile format using the Data Interoperability extension to create 4 output formats for each file. 10. Quality Assurance and rechecking of delivery files. The High Definition version is similar to how the layer exists in Landonline with a couple of changes to fix topology errors identified in topology checking. The following quality checks and steps were applied to the meshblock pattern: Translation of ESRI Shapefiles to ESRI geodatabase dataset The meshblock dataset was imported into the ESRI File Geodatabase format, required to run the ESRI topology checks. Topology rules were set for each of the layers. Topology Checks A tolerance of 0.1 cm was applied to the data, which meant that the topology engine validating the data saw any vertex closer than this distance as the same location. A default topology rule of "Must Be Larger than Cluster Tolerance" is applied to all data – this would highlight where any features with a width less than 0.1cm exist. No errors were found for this rule. Three additional topology rules were applied specifically within each of the layers in the ESRI geodatabase – namely "Must Not Overlap", "Must Not Have Gaps" and ""Area Boundary Must Be Covered By Boundary Of (Meshblock)". These check that a layer forms a continuous coverage over a surface, that any given point on that surface is only assigned to a single category, and that the dissolved boundaries are identical to the parent meshblock boundaries. Topology Checks Results: There were no errors in either the gap or overlap checks. Generalising To create the generalised Meshblock layer the "Simplify Polygon" geoprocessing tool was used in ArcGIS, with the following parameters: Simplification Algorithm: POINT\_REMOVE Maximum Allowable Offset: 1 metre Minimum Area: 1 square metre Handling Topological Errors: RESOLVE\_ERRORS Clipping of Layers to Coastline The processed feature class was then clipped to the coastline. The coastline was defined as features within the supplied Land2013 with codes and descriptions as follows: 11- Island – Included 12- Mainland – Included 21- Inland Water – Included 22- Inlet – Excluded 23- Oceanic – Excluded 33- Other – Included. Features were clipped using the Data

Interoperability extension, attribute filter tool. The attribute filter was used on both the generalised and high definition meshblock datasets creating four meshblock layers. Each meshblock dataset also contained all higher geographies and land-water data as attributes. Note: Meshblock 0017001 which is classified as island, was excluded from the clipped meshblock layers, as most of this meshblock is oceanic. Dissolve meshblocks to higher geographies Statistics New Zealand then dissolved the ESRI meshblock feature classes to the higher geographies, for both the full and clipped dataset, generalised and high definition datasets. To dissolve the higher geographies, a model was built using the dissolver, aggregator and sorter tools, with each output set to include geography code and names within the Data Interoperability extension. Export to MapInfo Format and Shapefiles The data was exported to MapInfo and Shapefile format using ESRI's Data Interoperability extension Translation tool. Quality Assurance and rechecking of delivery files The feature counts of all files were checked to ensure all layers had the correct number of features. This included checking that all multipart features had translated correctly in the new file.

## Metadata Constraints

### Constraints

#### Use Limitation

These conditions of supply apply to all users of Statistics NZ digital boundaries effective 1 July 2007. Permitted uses You must acknowledge Statistics NZ as the source of the boundaries. Uses not permitted You must not change the accuracy of the boundaries and supply them to another party. Liability While care has been taken to compile these boundary coordinates, Statistics NZ gives no warranty that the data supplied is free from error. Statistics NZ will not be liable for any loss suffered by the use, directly or indirectly, of this information.

## Metadata Constraints

### Legal Constraints

#### Use Limitation

Attribution 3.0 New Zealand

#### Use Limitation

<http://creativecommons.org/licenses/by/3.0/nz/>

#### Use Constraints

##### Restriction Code

license