

Metadata

File Identifier

585fb636-527e-acb0-9123-02ebc3c315e9

Language

eng

Hierarchy Level Name

dataset

Contact

Responsible Party

Individual Name

Geospatial Team

Organisation Name

Statistics New Zealand

Position Name

Geospatial Analyst

Contact Info

Contact

Phone

Telephone

Voice

04 931 4600

Address

Address

Delivery Point

Statistics House, The Boulevard, Harbour Quays

City

Wellington

Postal Code

6140

Country

NZ

Electronic Mail Address

Date Stamp**Date Time**

20161020

Metadata Standard Name

ISO 19115:2003/19139

Metadata Standard Version

1.0

Spatial Representation Info**Vector Spatial Representation****Integer**

0

Identification Info**Data Identification****Citation****Citation****Title**

REGC2013_V1_00

Abstract

This is the definitive set of regional council boundaries for 2013 as defined by the Local Government Commission and/or regional councils themselves but maintained by Statistics New Zealand (the custodian) as at 1 January 2013. The regional council is the top tier of local government in New Zealand. There are 16 regional councils in New Zealand (defined by Part 1 of Schedule 2 of the Local Government Act 2002). Eleven are governed by an elected regional council, while five are governed by territorial authorities (the second tier of local government) who also perform the functions of a regional council and are known as unitary authorities. These unitary authorities are Auckland Council, Nelson City Council, and Gisborne, Tasman, and Marlborough District Councils. The Chatham Islands Council also performs some of the functions of a regional council, but is not strictly a unitary authority. Unitary authorities act as regional councils for the purposes of a wide range of legislative purposes. Regional councils are responsible for the administration of many environmental and transport matters such as land transport planning and harbour navigation and safety. Regional Councils were established in 1989 after the abolition of the 22 local government regions. The Local Government Act 2002 requires the boundaries of regions to conform, as far as possible, to one or more water catchments. When determining regional boundaries, the Local Government Commission gave consideration to regional communities of interest when selecting which water catchments to include in a region. It also considered factors such as natural resource management, land use planning and environmental matters. Some regional council boundaries are coterminous with territorial authority boundaries but there are several exceptions. An example is Taupo District, which is geographically split between four regions, although most of its area falls within the Waikato Region. Where territorial local authorities straddle regional council boundaries, the affected

area is statistically defined by complete area units. In general, however, regional councils contain complete territorial authorities. The unitary authority of the Auckland Council was formed in 2010, under the Local Government (Tamaki Makarau Reorganisation) Act 2009, replacing the Auckland Regional Council and seven territorial authorities. The seaward boundary of any coastal regional council is the twelve mile New Zealand territorial limit.

Purpose

REGC2013_V1_00 is the definitive set of regional council boundaries for 2013. This version contains 16 regional councils.

Point Of Contact

Responsible Party

Individual Name

Geospatial Team

Organisation Name

Statistics New Zealand

Position Name

Geospatial Analyst

Contact Info

Contact

Phone

Telephone

Voice

04 931 4600

Address

Address

Delivery Point

Statistics House, The Boulevard, Harbour Quays

City

Wellington

Postal Code

6140

Country

NZ

Electronic Mail Address

geography@stats.govt.nz

Resource Constraints

Constraints

Use Limitation

<DIV STYLE="text-align:Left;"><DIV><DIV><P>These conditions of supply apply to all users of Statistics NZ digital boundaries effective 1 July

2007. Permitted uses You must acknowledge Statistics NZ as the source of the boundaries. Uses not permitted You must not change the accuracy of the boundaries and supply them to another party. Liability While care has been taken to compile these boundary coordinates, Statistics NZ gives no warranty that the data supplied is free from error. Statistics NZ will not be liable for any loss suffered by the use, directly or indirectly, of this information.

Language

eng

Topic Category Code

boundaries

Microsoft Windows 7 Version 6.1 (Build 7601) Service Pack 1; Esri ArcGIS 10.1.1.3143

Extent

EX _ Extent

Description

Data represents regional council polygons dissolved from meshblocks since 1991.

Extent

EX _ Extent

Geographic Element

EX _ Geographic Bounding Box

165.973643757-175.379047054-47.6201235791-33.9584971002

Distribution Info

Distribution

Transfer Options

Digital Transfer Options

On Line

Online Resource

Linkage

URL

<https://datafinder.stats.govt.nz/layer/25738-regional-council-2013/>

Data Quality Info

DQ _ Data Quality

Lineage

LI _ Lineage

Statement

Regional council boundaries are based on the meshblock pattern. Non-alignment of meshblock and cadastral boundaries are one of a number of reasons for meshblock boundary adjustments. Other reasons include requests from local authorities, Local Government Commission, Electoral Representation Commission and to make Census of Population and Dwellings enumeration processes easier. From the meshblock pattern, higher geographies, including the 2013 Regional council pattern were dissolved using the dissolve tool in the Arc GIS suite to create multiple output datasets.

Source

LI _ Source

Description

The original vertices delineating the meshblock boundary pattern were digitised in 1991 from 1:5,000 scale urban maps and 1:50,000 scale rural maps. The magnitude of error of the original digital points would have been in the range of +/- 10 metres in urban areas and +/- 25 metres in rural areas. Where meshblock boundaries coincide with cadastral boundaries the magnitude of error will be within the range of 1–5 metres in urban areas and 5 - 20 metres in rural areas. This being the estimated magnitude of error of Landonline. The creation of high definition and generalised meshblock boundaries for the 2013 digital pattern and the dissolving of these meshblocks into other geographies/boundaries were completed within Statistics New Zealand using ESRI's ArcGIS desktop suite and the Data Interoperability extension with the following process: 1. Import data and all attribute fields into an ESRI File Geodatabase from LINZ as a shapefile 2. Run geometry checks and repairs. 3. Run Topology Checks on all data (Must Not Have Gaps, Must Not Overlap), detailed below. 4. Generalise the meshblock layers to a 1m tolerance to create generalised dataset. 5. Clip the high definition and generalised meshblock layers to the coastline using land water codes. 6. Dissolve all four meshblock datasets (clipped and unclipped, for both generalised and high definition versions) to higher geographies to create the following output data layers: Area Unit, Territorial Authorities, Regional Council, Urban Areas, Community Boards, Territorial Authority Subdivisions, Wards, Constituencies and Maori Constituencies for the four datasets. 7. Complete a frequency analysis to determine that each code only has a single record. 8. Re-run topology checks for overlaps and gaps. 9. Export all created datasets into MapInfo and Shapefile format using the Data Interoperability extension to create 4 output formats for each file. 10. Quality Assurance and rechecking of delivery files. The High Definition version is similar to how the layer exists in Landonline with a couple of changes to fix topology errors identified in topology checking. The following quality checks and steps were applied to the meshblock pattern: Translation of ESRI Shapefiles to ESRI geodatabase dataset The meshblock dataset was imported into the ESRI File Geodatabase format, required to run the ESRI topology checks. Topology rules were set for each of the layers. Topology Checks A tolerance of 0.1 cm was applied to the data, which meant that the topology engine validating the data saw any vertex closer than this distance as the same location. A default topology rule of "Must Be Larger than Cluster Tolerance" is applied to all data – this would highlight where any features with a width less than 0.1cm exist. No errors were found for this rule. Three additional topology rules were applied specifically within each of the layers in the ESRI geodatabase – namely "Must Not Overlap", "Must Not Have Gaps" and ""Area Boundary Must Be Covered By Boundary Of (Meshblock)". These check that a layer forms a continuous coverage over a surface, that any given point

on that surface is only assigned to a single category, and that the dissolved boundaries are identical to the parent meshblock boundaries. Topology Checks Results: There were no errors in either the gap or overlap checks. Generalising To create the generalised Meshblock layer the “Simplify Polygon” geoprocessing tool was used in ArcGIS, with the following parameters: Simplification Algorithm: POINT_REMOVE Maximum Allowable Offset: 1 metre Minimum Area: 1 square metre Handling Topological Errors: RESOLVE_ERRORS Clipping of Layers to Coastline The processed feature class was then clipped to the coastline. The coastline was defined as features within the supplied Land2013 with codes and descriptions as follows: 11- Island – Included 12- Mainland – Included 21- Inland Water – Included 22- Inlet – Excluded 23- Oceanic – Excluded 33- Other – Included. Features were clipped using the Data Interoperability extension, attribute filter tool. The attribute filter was used on both the generalised and high definition meshblock datasets creating four meshblock layers. Each meshblock dataset also contained all higher geographies and land-water data as attributes. Note: Meshblock 0017001 which is classified as island, was excluded from the clipped meshblock layers, as most of this meshblock is oceanic. Dissolve meshblocks to higher geographies Statistics New Zealand then dissolved the ESRI meshblock feature classes to the higher geographies, for both the full and clipped dataset, generalised and high definition datasets. To dissolve the higher geographies, a model was built using the dissolver, aggregator and sorter tools, with each output set to include geography code and names within the Data Interoperability extension. Export to MapInfo Format and Shapefiles The data was exported to MapInfo and Shapefile format using ESRI's Data Interoperability extension Translation tool. Quality Assurance and rechecking of delivery files The feature counts of all files were checked to ensure all layers had the correct number of features. This included checking that all multipart features had translated correctly in the new file.

Metadata Constraints

Constraints

Use Limitation

These conditions of supply apply to all users of Statistics NZ digital boundaries effective 1 July 2007. Permitted uses You must acknowledge Statistics NZ as the source of the boundaries. Uses not permitted You must not change the accuracy of the boundaries and supply them to another party. Liability While care has been taken to compile these boundary coordinates, Statistics NZ gives no warranty that the data supplied is free from error. Statistics NZ will not be liable for any loss suffered by the use, directly or indirectly, of this information.

Metadata Constraints

Legal Constraints

Use Limitation

Attribution 3.0 New Zealand

Use Limitation

<http://creativecommons.org/licenses/by/3.0/nz/>

Use Constraints
Restriction Code
license